

Pharmacy Technician Certification Exam Answers

Section: Pharmacy Technician Certification Exams • Total Questions: 10 • Format: Verified Q&A

1. A physician has ordered heparin 7000 units subcutaneously daily. The medication is supplied in heparin 10,000 units in 1 mL. How many mL should be administered daily?

- (A) 0.07
 - (B) 0.14
 - (C) 0.7
 - (D) 1.42
-

2. Behind the counter decongestant products containing pseudoephedrine must be used with caution in patients with:

- (A) asthma
 - (B) hypertension
 - (C) hypokalemia
 - (D) eczema
-

3. Assuming proper storage conditions, a manufacturer expiration date of 09/26 means that the product's quality and efficacy is guaranteed until:

- (A) August 31, 2026.
 - (B) September 1, 2026.
 - (C) September 26, 2026.
 - (D) September 30, 2026.
-

4. A pharmacy technician receives a prescription for guaifenesin DM 1 tsp PO q.i.d. x 10 days. The technician prints a label that reads: Guaifenesin Take 10 mL by mouth two times a day for 10 days. What is the total number of errors this label contains?

- (A) 1
 - (B) 2
 - (C) 3
 - (D) 4
-

5. Rivastigmine has the same mechanism of action as:

- (A) Dorzolamide
 - (B) Doxazosin
 - (C) Donepezil
 - (D) Doxycycline
-

6. A pharmacy technician must provide a patient package insert to a patient who is picking up a prescription for which of the following types of medication?

- (A) Antibiotics
 - (B) Estrogen products
 - (C) Infused chemotherapy
 - (D) Antidiabetics
-

7. According to federal law, the information required on the prescription label for a controlled substance dispensed by the pharmacist in a community pharmacy shall contain the:

- (A) name and address of the prescriber
 - (B) pharmacy and prescriber's DEA registration numbers
 - (C) name of the patient and the name of the prescriber
 - (D) name and age of the patient
-

8. Once reconstituted, ciprofloxacin oral suspension stored in a refrigerator remains stable for a maximum of how many days?

- (A) 3
 - (B) 7
 - (C) 14
 - (D) 28
-

9. According to the Institute for Safe Medication Practices (ISMP) which of the following medications is considered a high-alert/high-risk medication?

- (A) Acetaminophen extra strength caplets
 - (B) Furosemide oral tablets
 - (C) Ceftriaxone sodium for injection
 - (D) Potassium chloride IV concentrate
-

10. Patients should avoid grapefruit juice while taking:

- (A) Simvastatin
- (B) Ibuprofen
- (C) Lisinopril
- (D) Doxycycline